

PORTARIA Nº 2505/2013

Concede Progressão na Carreira por Merecimento a Servidora Municipal.

O PREFEITO MUNICIPAL DE UMUARAMA, ESTADO DO PARANÁ, no uso de suas atribuições legais,

CONSIDERANDO o que estabelece os artigos 9 e 11, inciso III da Lei Complementar nº. 188/2007 de 19 de novembro de 2007, alterada pela Lei Complementar nº. 194/2007 de 21 de dezembro de 2007,

CONSIDERANDO ainda, o anexo VIII da já citada Lei, que estabelece a Tabela de Vencimentos/Salários;

RESOLVE:


Art. 1º. Conceder Progressão por Merecimento a Servidora Municipal **MARIA APARECIDA LOPES DA SILVA**, portadora da cédula de identidade RG nº 4.899.174-2 SSP/PR e inscrita no CPF nº 668.943.609-87, ocupante do Cargo de Carreira de Agente de Saúde, admitida em 20.03.2001 pelo regime Celetista, lotada no Fundo Municipal de Saúde, conforme estabelece os artigos 9 e 11, inciso III e anexo VIII da Lei Complementar nº.188/2007 de 19 de novembro de 2007, referente ao período de 20 de março de 2011 a 19 de março de 2013 passando da referência 12 para referência 13, classe "A", Grupo Ocupacional Administrativo I (GOA I).

Art. 2º. Esta portaria entra em vigor na data de sua publicação, produzindo seus efeitos a contar de 20 de março de 2013.

PAÇO MUNICIPAL, aos 17 de outubro de 2013.

MOACIR SILVA
Prefeito Municipal

ARMANDO CORDTS FILHO
Secretário de Administração



6. 1990. *Journal of the American Water Resources Association*, 26: 101-110.

—The *Journal of the American Medical Association* (JAMA) has published a study that suggests that the use of a single, low-dose, long-acting antipsychotic may be more effective than a combination of two or more antipsychotics in the treatment of schizophrenia. The study, conducted by researchers at the University of California, San Diego, found that patients who received a single, low-dose, long-acting antipsychotic (haloperidol decanoate) had a higher rate of response (75%) compared to those who received a combination of two or more antipsychotics (50%). The study also found that the single, low-dose, long-acting antipsychotic was associated with fewer side effects than the combination of two or more antipsychotics. The researchers conclude that the use of a single, low-dose, long-acting antipsychotic may be a more effective and safer treatment for schizophrenia than the combination of two or more antipsychotics.

6. Water, 100 ml was added to the flask and the mixture was stirred.

[illegible]

1. The first step is to identify the problem. This involves understanding the current situation and the goals that need to be achieved.

[illegible]

1. The first step in the process of the investigation is the identification of the problem. This is done by the investigator who is responsible for the investigation. The investigator must identify the problem and the scope of the investigation. The investigator must also identify the objectives of the investigation and the methods to be used. The investigator must also identify the resources available for the investigation.

[illegible]

Journal of Management Education 30(6)

PUBLICADO NO UMUARAMA ILUSTRADO
DE 20 / Outubro / 2013
DE Nº 9904
UMUARAMA, 21 / 10 / 2013
[Signature]
DIVISÃO DE ATOS OFICIAIS