



PREFEITURA MUNICIPAL DE UMUARAMA
ESTADO DO PARANÁ



PORTARIA Nº 325/2011

Prorroga prazo de validade da Portaria nº. 208, de 03 de fevereiro de 2011, que Constitui Comissão Especial.

O PREFEITO MUNICIPAL DE UMUARAMA, ESTADO DO PARANÁ, no
uso de suas atribuições legais,

RESOLVE:


Art. 1º. Fica prorrogado o prazo da Portaria nº. 208, de 03 de fevereiro de 2011, que Instaurou Processo Administrativo Disciplinar, com fundamento no artigo 152 da Lei Complementar nº. 018/1992, com o objetivo de apurar aptidão da servidora, lotada na Secretaria Municipal de Educação, por mais 30 (trinta) dias.

Art. 2º. Esta Portaria entra em vigor a partir da data de sua publicação.

PAÇO MUNICIPAL, aos 28 de fevereiro de 2011.

MOACIR SILVA
Prefeito Municipal

ARMANDO CORDTS FILHO
Secretário de Administração



ESTADO DE PARANÁ
PREFEITURA MUNICIPAL DE MARACÁI



1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1972). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1972). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total moisture content was determined by the method of AOAC (1990). The total dry matter content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenolic content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenolic content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).

the 1990s, the number of people in the United States who are 65 years of age or older is projected to increase from 20 million to 30 million, and the number of people 75 years of age or older is projected to increase from 10 million to 15 million (U.S. Census Bureau, 1996).

[illegible][illegible]

1. The first step is to identify the problem. This involves understanding the current situation and the goals that need to be achieved.

the "good" and "bad" of the "new" and "old" is a matter of perspective.

Journal of Management Studies, 19(1), 67-80.

PUBLICADO NO UMUARAMA ILUSTRADO
DE 02 | março | 20 11
DE Nº 9.118
UMUARAMA, 02 | 03 | 20 11
DIVISÃO DE SERVIÇOS GERAIS E PATRIMÔNIO